

HR1156 LRB098 21519 GRL 60086 r

1 HOUSE RESOLUTION WHEREAS, 1 in 150 children in the United States are born 2 3 with congenital cytomegalovirus; and WHEREAS, Congenital cytomegalovirus is the most common 4 cause of birth defects and childhood disabilities in the United 5 6 States; and 7 WHEREAS, Congenital cytomegalovirus is preventable with 8 behavioral interventions, such as practicing frequent hand 9 washing with soap and water after contact with diapers or oral 10 secretions, not kissing young children on the mouth, and not sharing food, towels, or utensils with young children; and 11 12 WHEREAS, Cytomegalovirus is found in bodily fluids, such as 13 saliva, blood, urine, and tears; and WHEREAS, Congenital cytomegalovirus can be diagnosed if 14 15 the virus is found in saliva, blood, or other body tissues of an infant during the first week after birth; and 16 17 WHEREAS, Most people are not aware of their cytomegalovirus infection status, with pregnant women being one of the highest 18 19 risk groups; and

- 1 WHEREAS, The American College of Obstetricians and
- 2 Gynecologists and the Centers for Disease Control and
- 3 Prevention recommend that OB/GYNs counsel women on basic
- 4 prevention measures to guard against cytomegalovirus
- 5 infection; and
- 6 WHEREAS, The incidence of children born with congenital
- 7 cytomegalovirus can be greatly reduced with public education
- 8 and awareness; and
- 9 WHEREAS, A comprehensive understanding of cytomegalovirus
- 10 provides opportunities to improve the health and well-being of
- 11 our children; therefore, be it
- 12 RESOLVED, BY THE HOUSE OF REPRESENTATIVES OF THE
- 13 NINETY-EIGHTH GENERAL ASSEMBLY OF THE STATE OF ILLINOIS, that
- we designate the month of June 2014 as Illinois Cytomegalovirus
- 15 Awareness Month in the State of Illinois in order to raise
- 16 awareness of the dangers of cytomegalovirus and reduce the
- 17 occurrence of congenital cytomegalovirus infection; and be it
- 18 further
- 19 RESOLVED, That we recommend that more effort be taken to
- 20 counsel women of childbearing age of the effect this virus can
- 21 have on their children.